
Short Answers and Questions

Q1: Why the element of group 1 are called alkali metals?

Answer

The elements of group 1 are called alkali metal because they produce alkalies with H_2O .

Q2: Describe the trends in atomic radius.

Answer

The number of shells in all the elements of a given periods remains the same but the value of effective nuclear charge, increases from left to right. The increased effective nuclear charge pulls the electron cloud of the atom nears to the nucleus and thus the size of the atoms and ions goes on decreasing from left to right. This in going from left to right in a period of S and P block elements atomic and ionic radii decreased with the increase of atomic number.

Q3: Describe the trends in electronegativity?

Answer

Electronegativity is a measure of tendency of an atom to attract a bonding pair of electrons. The pauling scale is the most commonly used. Fluorine (the most electronegative element) is assigned a value of 4.0, and values range down caesium and francium which are least electronegative at 0.7.

- =====
- 1) On moving from left to right in a period, there is a decrease in the size of atoms, smaller atoms have greater tendency to attract the electrons towards themselves. Smaller atoms have higher electronegativity values.
 - 2) On a moving from left to right in a period there is an increase of ionization energy and electron affinity of the elements. The atoms of the elements which have higher value of ionization energies and electron affinities also have higher electronegativities.

Q4: Explain that sodium magnesium and aluminium all have metallic structure.

Answer

In sodium only one electron per atom is involved in the metallic bond the single 3s electron. In magnesium both of its outer electrons are involved and in aluminium in all three. Sodium is 8 – coordinated each sodium atom is touched by only 8 other atoms.

Both magnesium and aluminium are 12 – coordinated. This is more efficient way to pack atoms, leading to less wasted space in the metal structure and to stronger bonding in the metal.

Q5: Silicon has a giant covalent structure just like diamond structure justify it.

Answer

Silicon has a giant covalent structure just like diamond. A tiny part of the structure is held together by strong covalent bonds in all three dimensions.

Q6: Sodium, magnesium and aluminium are good conductors of electricity? Why?

Answer

Sodium, magnesium and aluminium are good conductors of electricity. Conductivity increases as you go from sodium to magnesium to aluminium as they have free electrons. Silicon is a semiconductor.

The 3 metals, conduct electricity because the delocalised electrons are free to move throughout solid or the liquid metal.

Q7: Phosphorus, sulphur, chlorine and argon have lower melting and boiling points than those of the first four members of the period which have giant structure justify it.

Answer

Phosphorus, sulphur, chlorine and argon are simple molecular substances with only van der Waals attraction between the molecules. The size of the melting and boiling point are governed by the sizes of the molecules.

Phosphorus: contains P_4 molecules. To melt phosphorus you don't have to break any covalent bond forces between the molecules.

Sulphur: Sulphur consists of S_8 rings of atoms. The molecules are bigger than phosphorus molecules and so the van der Waals attraction will be stronger, leading to a higher melting and boiling point.

Chlorine: Chlorine Cl_2 is a much smaller molecule with comparatively weak van der Waals attraction and so chlorine will have a lower melting and boiling point than sulphur or phosphorus.

Argon: Argon molecules are just single argon atom, Ar. The scope for van der Waals attractions between these is very limited and so the melting and boiling of argon are lower again.

Q8: Write down the chemical reaction of the period 3 element with oxygen and chlorine?

Answer


Reaction with oxygen

Sodium

For the simple oxide.



For peroxide



Magnesium:

Mg burns in oxygen with an intense white flame to give white solid magnesium oxide.



Silicon:

Silicon will burn in oxygen if heated strongly.

Silicon dioxide is produced.



Phosphorus:

For the phosphorus (III) oxide.



For phosphorus (V) oxide.



Sulphur:

Sulphur burns in air or oxygen or gentle heating with a pale blue flame.



Chlorine and argon:

Despite having several oxides, chlorine won't react directly with oxygen.

Argon doesn't react either.

Q9: Write down the reaction of Aluminium and phosphorus with chlorine?

Answer

Aluminium is often reacted with chlorine by passing dry chlorine over aluminium foil heated in a long tube. The aluminium burns in the stream of chlorine to produce very pale yellow aluminium chloride. This sublimes and collects further down the tube where it is cooler.



White phosphorus burns in chlorine to produce a mixture of two chlorides, phosphorus

(III) chloride and phosphorus (V) chloride (phosphorus trichloride and phosphorus pentachloride.

Phosphorus (III) chloride is a colourless fuming liquid.



Phosphorus (V) chloride is an off white (going towards yellow) solid.


Q10: Write down the reaction of silicon and sulphur with chlorine.

Answer

When chlorine is passed over silicon powder heated in a tube, it reacts to produce silicon tetrachloride. This is a colourless liquid which vaporises and can be condensed further along the apparatus.



Sulphur: When a stream of chlorine is passes over some heated sulphur, it reacts to form an orange, will smelling liquid disulphur dichloride S_2Cl_2 .



Q11: Write down physical properties of Phosphorus oxides.

Answer

Phosphorus Oxide has tow common oxides

- 1) Phosphorus (III) oxide. P_4O_6
- 2) Phosphorus (V) oxide P_4O_{10}

Phosphorus oxides

Phosphorus (III) oxide is a white solid, melting at 24°C and boiling at 173°C . The structure of its molecule it best worked out stating from a P_4 molecules which is a little tetrahedron.



P_4 molecule

The phosphorus is using only 3 of its outer electrons (the 3 unpaired P electrons) to forms bonds with the oxygens.

Phosphorus (V) oxide

Phosphorus oxide is also a white solid, subliming at 300°C . In this case the phosphorus uses all five of its outer electrons in the bonding.

Solid phosphorus (V) oxide exists in several different forms some of them polymeric. The force oxygens are attached to the faces phosphorus atoms in a double bonds.


Q12: Write down the Physical properties of sulphur oxide.

Answer

Sulphur has two common oxides, sulphur dioxide SO_2 and sulphur trioxide SO_3 .

Sulphur dioxide

Sulphur dioxide is a colourless gas at room temperature with an easily choking smell. It consists of SO_2 molecules.



The sulphur uses 4 of its outer electrons to form the double bond with the oxygen leaving the outer two as a lone pair on the sulphur. The bent shape of SO_2 is due to this lone pair.

Sulphur trioxide

Sulphur trioxide is a white solid with a low melting point. It consists of simple SO_3 molecules in which all six of the sulphur's outer electrons are involved in the bonding.



There are various forms of solid sulphur trioxide. The simplest one is a trimer S_3O_9 where 3 SO_3 molecules are joined up and arranged in a ring.

There are also polymeric forms in which the SO_3 molecules join together in long chains.



The simple molecules join up in this way to make bigger structures. This is what makes the sulphur trioxide a solid rather than a gas.


Q13: Write down the physical properties and structures of chlorine oxides.

Answer

Chlorine forms several oxides: chlorine (I) oxide Cl_2O and chlorine (VII) oxide.

Chlorine (I) oxide

Chlorine (I) oxide is a yellowish red gas at room temperature. It consists of simple small molecules.




In this structure chlorine uses its one outer electron and bonds with oxygen.

Chlorine (VII) oxide

Chlorine (VII) oxide is a colourless oily liquid at room temperature.

In chlorine (VII) oxide the chlorine uses all of its seven outer electrons and bonds with oxygen. This produces a much bigger molecule.



Q14: Why the melting and boiling points of metal oxide and silicon dioxide is high?

Answer

The metal oxides and silicon dioxide (the giant structures) will have high melting and boiling point because a lot of energy is needed to break the strong bonds operating in 3 dimensions.

The attractive forces between the oxides of phosphorus, sulphur and chlorine will be van der Waals dispersion and dipole-dipole interaction. These vary in size depending on the size, shape and polarity of the various molecules but will always be much weaker than the ionic or covalent bonds need to break in a giant structure.


Q15: Write down the reaction of sodium oxide (Na_2O) with acids and H_2O .

Answer

Sodium oxide (Na_2O) is a simple strongly basic oxide. It is basic because it contains the oxide ion, O^{2-} which is a very strong base with a high tendency to combine with hydrogen ions.

Reaction with H_2O

Na_2O reacts exothermically with cold H_2O to produce sodium hydroxide solution.



Reaction with acids



Q16: Write down the reaction of Magnesium Oxide (MgO) with H_2O and acids.

Answer

MgO is a simple basic oxide because it also contains oxide ions. In Magnesium oxide case the attractions are between $2+$ and $2-$ it takes more energy to break these.

Reaction with H_2O

If you shake some white magnesium oxide with H_2O it doesn't look as if it reacts. If you test the pH of the liquid, you find that it is somewhere around pH 9. There must be some slight reaction with H_2O to produce hydroxide ions in solution. Some $\text{Mg}(\text{OH})_2$ is formed in the reaction, but this is almost insoluble and so not many hydroxide ions actually get into solution.

Reaction with acid



MgO reacts with acid to form MgCl_2 solution.




Q17: Write down the reaction of phosphorus (V) oxide (P_4O_{10}) with H_2O and Base?

Answer

Phosphorus (V) oxide reacts violently with H_2O to give a solution containing a mixture of acids, the nature of which depends on the conditions, use usually ust consider one of these, phosphoric acid H_3PO_4 (also known just as phosphoric acid or as orthophosphoric acid).



Reactions with Base



Again, if you were to react phosphorus (V) oxide directly with sodium hydroxide solution rather than making the acid first you would end up with the same possible salts.

This is getting ridiculers, and so will only give one example out of the possible equations.



Q18: Write down the reaction of sulphur dioxides (SO_2) with H_2O and base?

Answer

Reaction with H_2O

Sulphur dioxide is fairly soluble in H_2O reacting which it to give a solution of sulphurs acid H_2SO_3 .



Reaction with Base

As it is acidic so it reacts with $NaOH$ and CaO as follows.



Q19: Write down the reaction of sulphur trioxide with Base?

Answer

Sulphur trioxide reacts violently with H_2O to produce a fog of concentrated sulphuric acid dioplets.



Reacts with Base



Sulphur trioxide itself will also react directly with bases to form sulphates. e.g. it will react with CaO to form calcium sulphates.



Q20: Write down the reaction of chlorine oxides (Cl_2O) with H_2O and Base.

Answer

Chlorine forms several oxides but the only two are chlorine (VII) oxide Cl_2O_7 and chlorine (I) oxide, Cl_2O , and chlorine (I) oxide, Cl_2O . Chlorine (VII) oxide is also known as dichlorine heptoxide and chlorine (I) oxide as dichlorine monoxide.

Chlorine (VII) oxide

Chlorine (VII) oxide is the highest oxide of chlorine the chlorine is in its maximum oxidation state of +7. It continues the trend of the highest oxides of the period 3 elements towards being stronger acids.

Reaction with H_2O

Chlorine (VII) oxide reacts with H_2O to give the very strong acid chloric (VII) acid also known as perchloric acid. The pH of typical solution will be around 0. The pH of typical solution will be around 0.



Reaction with Base

Chloric (VII) acid reacts with NaOH solution to form a solution of sodium chlorate (VII).




Chlorine (VII) oxide itself also reacts with sodium hydroxide solution to give the same product.



Q21: Write down the reaction of chlorine (I) oxide with Base?

Answer

Chlorine (I) oxide is far less acidic than chlorine (VII) oxide. It reacts with water to some extent to give chloric (I) acid, HOCl also known as hypochlorous acid.



Reaction with Base

Chloric (I) acid reacts with sodium hydroxide solution to give a solution of sodium chlorate (I) (sodium hypochlorite).



Chlorine (I) oxide also reacts directly with sodium hydroxide to give the same product.



Q22: Give a detail note on sodium chloride NaCl.

Answer

Sodium Chloride is a simple ionic compound consisting of a giant array of sodium and chloride ions.

The strong attractions between the positive and negative ions need a lot of heat energy to break, and so sodium chloride has high melting and boiling points.

It doesn't conduct electricity in the solid state because it hasn't any mobile electrons and the ions aren't free to move. However, when it melts it undergoes electrolysis.

Sodium chloride simply dissolves in H_2O to give a neutral solution.

Q23: Give a detail note on Magnesium chloride $MgCl_2$.

Answer


Magnesium chloride is also ionic, but with a more complicated arrangement of the ions to allow for having twice as many chloride ions as magnesium ions.

Again, lots of heat energy is needed to overcome the attractions between the ions, and so the melting and boiling points are again high.

Solid magnesium chloride is a non-conductor of electricity because the ions aren't free to move. However, it undergoes electrolysis when the ions become free on melting.

$MgCl_2$ dissolves in H_2O to give a faintly acidic solution (pH = approximately 6). When magnesium ions are broken off the solid lattice and go into solution, there is enough action between the $2+$ ions and the water molecules to get co-ordinate bonds formed between the magnesium ions and lone pairs on surrounding H_2O molecules.

Hexaquaamagnesium ions are formed $[Mg(H_2O)_6]^{2+}$



Q24: Give a detail note on Aluminium chloride $AlCl_3$.

Answer


At room temperature, solid aluminium chloride has an ionic lattice with a lot of covalent character.

At temperature around $180 - 190^\circ C$ (depending on the pressure) aluminium chloride connects to a molecular form Al_2Cl_6 . This causes it to melt or vaporise because there are now only comparatively weak inter-molecular attractions. As the temperature increases a bit more, it increasingly breaks up into simple $AlCl_3$ molecules.

Solid aluminium chloride doesn't conduct electricity at room temperature because the ions are not free to move.

Molten aluminium chloride doesn't conduct electricity because there are not any ions any more.

The aluminium chloride reacts with H_2O rather than just dissolving in it. In the first instance, hexaquaaluminium ions are formed together with chloride ions.



Q25: Write down the details note on silicon tetrachloride SiCl_4 .

Answer

SiCl_4 is a simple covalent chloride between the silicon and the chlorine for the two to form ionic bonds.

Silicon tetrachloride is a colourless liquid at room temperature which fumes in moist air. The only attractions between the molecules are van der Waals dispersion forces.

It doesn't conduct electricity because of the lack of ions mobile electrons.

It fumes in moist air because it reacts with H_2O in the air to produce hydrogen chloride. If you add H_2O to silicon tetrachloride there is a violent reaction to produce silicon dioxide and fumes of hydrogen chloride.

In a large excess of H_2O the hydrogen chloride will of course, dissolve to give a strongly acidic solution containing hydrochloric acid.



Q26: Write down the reaction of phosphorus chlorides with H_2O .

Answer


There are two phosphorus chlorides phosphorus (III) chloride, PCl_3 and phosphorus (V) chloride PCl_5 . Phosphorus (III) chloride (phosphorus trichloride) PCl_3 .

This is simple covalent chloride again a fuming liquid at room temperature. Phosphorus (III) chloride reacts violently with H_2O .



Phosphorus (V) chloride (Phosphorus pentachloride) PCl_5 .

Phosphorus (V) chloride is a white solid which sublimes at 163°C . The higher the temperature goes above that, the more the phosphorus (V) chloride dissociates to give phosphorus (III) chloride and chlorine.



Solid phosphorus (V) chloride contains ions that is a solid at room temperature. The formation of ions involves two molecules of PCl_5 .

Phosphorus Chloride has a violent reaction with H_2O producing fumes of hydrogen chloride.

As with the other covalent chlorides, if there is enough H_2O present, these will dissolve to give a solution containing hydrochloric acid.



If H_2O is boiling, the phosphorus (V) chloride reacts further to give phosphorus (V) acid and more HCl . Phosphoric (V) acid is also known just as phosphoric acid or as orthophosphoric acid.



The overall equation in boiling H_2O is just a combination of these.



=====

Q27: Write down the formula of following:

1) Ortho silicic acid


3) Sulphuric acid

2) Phosphoric acid


4) Chloric acid

Answer

Orthosilicic acid.




Phosphoric acid



Sulphuric acid



Chloric acid




Q28: Aluminium hydroxide is amphoteric. Justify it.

Answer

Aluminium hydroxide is amphoteric like Na and Magnesium hydroxides, it will react with acids. This is showing the basic side of its nature.

With dilute hydrochloric acid, a colourless solution of aluminium chloride is formed.



But aluminium hydroxide also has an acidic side to its nature. It will react with sodium

=====

hydroxide solution to give a colourless solution of sodium tetrahydro aluminate.



Q29: Why atomic Radii increases as we move from Li to caesium?


Answer

As we move from lithium to caesium an extra shell of electrons of an extra shell increases there in an in increases of atomic and ionic radii (of M^+ cons) as we move from lithium to caesium.

Q30: Define ionization energy and why Alkali metals have low ionization energy?

Answer

First ionization energy is the energy needed to remove the most loosely held electron from each of one mole of gaseous atoms to make one mole of singly charged gaseous ions, in other words for 1 mole of this process.



As you go down the graph ionization energy falls. Alkali metal have only one electron in their outermost shell (ns1 electron). The ns1 electron is so weakly held with the nucleus that it can be removed very easily. Alkali metals therefore have low ionisation energies.

As the distance ns1 electron from the nucleus increases on moving from Li to Cs its removal becomes more and more easy as are proceed from Li to Cs i.e the amount of energy used in the removal of ns1 electron is maximum in case of energies of alkali metals go on decreasing from Li to Cs.

The second ionisation energies are fairly high, since the loss of the second electron from M^+ cation which has a noble gas configuration is quite difficult.

Q31: Define electronegativity. And write down its trend in alkali metals.

Answer

Electronegativity is a measure of the tendency of an atom to attract a bonding pair of electrons. In alkali metals to the outer electron of the atom of alkali metals is loosely held with the nucleus and hence it can be easily excited to the higher energy levels even by a small amount of heat energy. During the excitation process the electron absorbs some energy and when this excited electron comes back to its original position, it gives out absorbed energy in the form of light in visible region of the electromagnetic. Since the amount of energy absorbed during the excitation process is different in different atoms, different colours are imparted by the atom to the flame. The property of alkali metal to give colouration in the burn flame has been used to detect their presence in salts by a test, known as flame test.

Q32: Why the melting and boiling point of group 1 elements go down the group?

Answer

The melting and boiling points are very low because of the presence of weak inner atomic bonds in the solid state of the alkali metals. These bonds are due to their atomic radii and mainly due to their electronic configuration having a single valence electrons are compared to large number of available vacant orbital.

As the size of the metal atoms increases the repulsion of non bonding electrons

decreases the melting and boiling points of alkali metals when we move from Li to Cs.

Q33: The densities of alkali metal are low, why?


Answer

The densities of alkali metals are low due to large atomic volumes. Li, Na, and K are lighter than H₂O. The densities increase with the increases in atomic from Li to Cs indicating the greater atomic weight more than compensates for the bigger size of the atoms. K is however lighter than Na which is due to an _____ increase in atomic size of K.


Q34: Write down the reaction of first group alkali metal with oxygen.

Answer

Alkali metals react with O₂ in air rapidly and thus tarnished due to the formation of their oxide on the surface of the metals. It is for this reason that alkali metals are stored in kerosene or paraffin oil.





Na when burnt in O₂ forms sodium peroxide Na₂O₂.




Other alkali metals react with O₂ to form superoxide of MO₂ type.





Normal oxides of alkali metals other than that of Li (Li₂O) are not formed by the direct reaction b/w the metals and O₂ they are formed by indirect methods by producing peroxides, nitrates, nitride with metals itself.



Normal oxides react with H₂O to form hydroxides by proton exchange.



The peroxides (O₂²⁻) and superoxides (O₂²⁻) are strong oxidising agents and react with H₂O to give H₂O₂ and O₂.



Q35: Write down effect of Nitrates, carbonates and hydrogen carbonates.


Answer

Group 1 compounds are more stable to heat than the corresponding compounds in group 2. Li compounds behave similarly to group 2 compounds but the rest of group 1 are in same way different.

Nature of carbonates, bicarbonates and Nitrates

The carbonates (M_2CO_3) and bicarbonates ($MHCO$) are highly stable to heat with increase of electropositive character from Li to Cs, the ability of these salts increases.

Their nitrates decompose on strong heating to the corresponding nitrate and O_2 .



Q36: Explaining the trend in terms of the polarising ability of the positive ion. Justify.

Answer

When alkali metal cations approach near an anion attracts the outer most electron of the anion and repels the molecules. Thus the distortion of polarisation of the anion takes place. This distortion usually showing of electron L/w two oppositely charged ions the bond between the cations and anion becomes partly covalent in character. In general the smaller cations polarise the anions more effectively than bigger one. Therefore, the lithium salts are slightly covalent which other alkali metal salts are ionic.

Q37: Define flame test? And the property of alkali metals to give also reaction in the Bunsen flame has been used to detect the presence in salts by a test known as flame test? Justify.

Answer

Flame test are used to identify the presence of a relatively small number of metal ions in a compound.

We have seen that outer electron of atom of alkali metals is loosely held with nucleus and hence it can be easily excited to the higher energy levels even by a small amount of heat energy. During the excitation process the electron absorbs some energy and under this excited electron comes back to its original position, it gives out absorbed energy in the form of light in visible region of the electromagnetic spectrum and hence the colour is imparted by the atoms to the flame. Since the amount of energy absorbed to the flame. The property of alkali metals to give coloration in the Bunsen flame has been used to detect their presence in salts by a test known as flame test.

Q38: Alkaline earth metals have higher nuclear charge which tends to draw the orbit electrons towards the nucleus?

Answer

Because of the addition of an extra shell of electrons to each element from Be to Ra, the atomic volume increases from Be to Ra. With the increase of atomic volume the atomic and ionic radii (of M^{2+} ions) also increases from Be to Ra. The atomic radii of these elements are however smaller than those of alkali metals have higher nuclear charge which tends to draw the electrons towards the nucleus. The smaller values of atomic radii result in that the alkaline earth metals are harder, have higher densities and higher melting points than alkali metals.

Q39: $Mg(OH)_2$ is weakly basic while $Be(OH)_2$ is amphoteric why?

Answer

$Be(OH)_2$ is not at all basic in fact it is amphoteric since it reacts with acids to form salts and with alkalis to give beryllates.



The hydroxides of other metals are basic character. Their basic character increase on moving down the group. Thus $\text{Mg}(\text{OH})_2$ is weakly basic while $\text{Be}(\text{OH})_2$ is the amphoteric. The increase in basic character of the hydroxides on moving down the group is due to the fact that with the increase in size of M^{2+} cation both the polarity of $\text{M}-\text{OH}$ bond and the internuclear distance between oxygen of OH ion and the metal atom increase. As a result of this there is greater ionisation of $\text{M}(\text{OH})_2$ and hence basic character increases.


Due to high polarising power of small Be^{2+} in $\text{Be}(\text{OH})_2$ are almost insoluble in H_2O while the hydroxides of other metals are slightly soluble. Their solubility increases on moving down the group as shown by the increasing value of the solubility products of these hydroxides.

Q40: Discuss the effects of heat on the group 2 nitrates.

Answer

All the nitrates in this group undergo thermal decomposition to give the metal oxide, nitrogen dioxide and oxygen.

The nitrates are white solids, and the oxides produced are also white solids. Brown nitrogen dioxide gas is given off together with oxygen. Magnesium and calcium nitrates normally have water of crystallization and the solid may dissolve in its own H_2O of crystallization to make a colourless solution before it starts to decompose.



As we go down the group, nitrates also have to be heated more strongly before they will decompose.

Q41: How Beryllium differs from members of its family?

Answer

Be is the first element of the group and differs from rest of alkaline earth metals due to its small atomic size and comparatively high electronegativity.

Hardness and Melting and boiling point

Be is hardest by all the elements of its group and their melting and boiling points are also high.

Formation of Covalent Bond

Be has a tendency to form covalent compounds. Thus when it reacts with other elements the electronegativity difference is small and the bond is therefore covalent.

Reaction with H_2O

Be does not react with H_2O even at high temperature, other alkaline earth metals decompose water liberating H_2 gas.



Rex with alkalis

Be react with alkalis to form hydrogen



Behaviour of oxides and hydroxides

The oxides and hydroxides of Be are amphoteric dissolve in both acid and alkalis to form salts.




Q42: Write down the number of molecules of H_2O of crystallization of Be^{2+} .

Answer

The salts of Be^{2+} ion cannot have more than four molecules of H_2O of crystallisation while other alkaline earth metals have more than four molecules of H_2O of crystallisation. This is explained as follow. In case of Be^{2+} ion there are only four orbitals (namely one orbitals can accept lone pairs of electrons slenobed by O – atoms on each of the H_2O molecules are as.


Valance shell electronic configuration

of Be atom = $2s^2, 2p^0$




Valance shell configuration of Be^{2+}

($2s^0, 2p^0$)



Attachment of $4\text{H}_2\text{O}$ molecules with

Be^{2+} ion



Other alkaline earth metals like Mg can ____ their coordinate ion no to six by using one their outer most shells.

Q43: The melting and boiling pointing C to Pb are gradually decreases down the group. Why?

Answer

As we move down the group from c to Pb the melting point as well as boiling points generally decreases, although the decrease is not in a regular order. This decrease in melting points as well as in boiling point indicates that inter-atomic forces also decreases in the same direction. The melting and boiling point of c and Si are notably high because of the tendencies of these elements to form giant molecules. The low value for tin's melting point compared with lead is presumably due to forming a distorted 12 co – ordinate structure rather than a pore one.

Q44: Carbon and silicon doesn't conduct electricity.

Answer

Carbon as diamond doesn't conduct electricity. In diamond the electrons are all tightly

bond and not free to move.

Unlike diamond (which doesn't conduct electricity) silicon, germanium and grey tin are semiconductor.

White tin and lead are normal conductors of electricity. There is clear trend from the typically non metallic conductivity behaviour of carbon as diamond, and the typically metallic behaviour of white tin and lead.

Q45: Electronegativity or ionization energy decreases of moving down type group from c to Pb. Why?

Answer

Carbon is the most electronegative element of this sub group and the electronegativities decreases with like of atomic no but not a regular manner. This is probably due to the filling of the d orbitals in case of Ge and Sn and f orbitals in case of Pb.

The ionisation energy values decrease on moving down the group from c to Pb although the decreases does not occur in a regular order. The irregulating in the decreases of those values is due to the filling of intervening d – orbitals in case of Ge and Sn and f – orbitals in case of Pb which are not able to seen the valance electrons effectively in elements.

Q46: Carbon and silicon show +4 oxidation state while occurrence of +2 and +4 oxidations states in case of Ge, Sn and Pb. Explain.

Answer


Carbon and silicon show + 4 oxidation state while occurrence of + 2 and +4 oxidation states in case of Ge, Sn and Pb are when only two np electrons from the ns^2np^2 configuration are lost. The element in +2 oxidation states remain inert and hence are not lost in the formation of M^{2+} cations. This pair of ns^2 electrons is called inter pair of electrons. Since the group the stability of +2 oxidation state also increases from Ge^{2+} to Pb^{2+} i.e. $Ge^{2+} < Sn^{2+} < Pb^{2+}$.

When all four ns^2np^2 electrons are lost we get the elements in +4 oxidation state i.e. M^{4+} cations are formed. On decreasing the group stability of +4 oxidations state decreases i.e. the stability of M^{4+} cations decreases from Ge^{4+} to Pb^{4+} . i.e. $Ge^{4+} > Sn^{4+} > Pb^{4+}$.

Q47: The dipole moment of CO_2 and explain it?

Answer

The dipole moment of CO_2 is zero. Therefore it is a linear molecule.



Q48: Draw and explain the structure of SiO_2 .

Answer

It is a macro molecular compounds in which silicon and oxygen atoms are linked together covalently in tetrahedral basic unit.


In crystablite, these unit are joined as in diamond, circle in quartz and tridymite they are arranged spirally around an axis. Because of its structure of SiO_2 is non volatile and hard unlike CO. Triatomic molecules of SiO_2 and CO_2 , carbon and silicon are similar

in having.

- 1) 4 valence electron
- 2) 4 covalent bond formation

But they show a lot of difference in their physical properties. It is due to the fact that.

- 1) Silicon atoms are much larger in size than C atoms and thus tend to be surrounded by more oxygen atoms.
- 2) Silicon forms only single bond with oxygen atom while carbon forms double bonds.



Si atom is bound to four oxygen atom in a tetrahedral structure which results in the formation of SiO_2 crystal. The simplest formula for silica is SiO_2 .

Q49: Write down the trends of electronegativity of Halogens and also its first ionization energy.

Answer

Halogens have high values of electronegativity in the group. These values decrease as they proceed from F to I in the group. Large electronegativity values of halogen atoms indicate that X atoms have a strong tendency to form X^- ions. Electron affinity values decrease from Cl to I.

Q50: Define bond enthalpies and also explain the bond enthalpies in halogens?

Answer

Bond enthalpy is the heat needed to break one mole of a covalent bond to produce individual atoms, starting from the original substance in the gas state, and ending with gaseous atoms.

The bond enthalpies of the $\text{Cl}-\text{Cl}$, $\text{Br}-\text{Br}$ and $\text{I}-\text{I}$ bonds fall just as you would expect, but the $\text{F}-\text{F}$ bond is deviated from the sequence. This is because due to very small $\text{F}-\text{F}$ bond length, there is a very large repulsion between the lone pairs of the F atoms in F_2 molecules, which helps the dissociation of F_2 molecules into atoms.

$\text{X}-\text{X}$ bonds in Cl_2 , Br_2 and I_2 molecules are stronger than $\text{F}-\text{F}$ bond in F_2 molecule. This is due to the possibility of the existence of multiple bonds in $\text{X}-\text{X}$ bonds involving d-orbitals.

Q51: HF is the weakest acid. Why?

Answer

HF ionises only slightly while HCl, HBr, and HI ionise completely. Hence HF is the weakest acid and the strength of these acids increases from HF to HI.

i.e. HF (weakest acid) < HCl < HBr < HI.

The weakest acidic nature of HF is due to the fact that the dissociation energy of H-F bonds in HF molecule is the highest and hence this molecule has least tendency to split up into H^+ and F^- ion in aqueous solution.

Q52: Define open Hearth process.

Answer

A fire furnace is used to remove the impurities of metals. It is lined inside with fire bricks and is just like a room. Burning gases are entered from one side and exhaust gases are removed from the opposite end. The process is operated from opposite ends after an interval. Metals melt in a shorter time by this two-way heating.

Q53: Write down the application of bleaching powder.

Answer

Bleaching powder is actually a mixture of calcium hypochlorite ($Ca(OCl)_2$) and the basic chloride $CaCl_2$. H_2O will form slaked lime $Ca(OH)_2$.

Bleaching powder is used for the disinfection of drinking H_2O or swimming pools H_2O . It is used as a sanitizer in outdoor swimming pools in combination with a cyanuric acid stabilizer which reduces the loss of chlorine due to ultraviolet radiation.

The calcium content hardens the H_2O and tends to clog up some filters hence some products containing calcium antiscaling agents.

Bleaching powder is used for bleaching cloth and linen. It is also used in bathroom, cleaners, household disinfectant sprays, moss and algae removers and cures/____.

In addition, bleaching powder may be used to manufacture chloroform. Bleaching powder is used also in sugar industry for bleaching sugar cane juice before its crystallization.

Q54: Define Goiter?

Answer

The term goiter refers to the abnormal enlargement of the thyroid gland due to deficiency of iodine in diet. It results in swelling in the neck. It is important to know that the presence of goiter does not necessarily mean that the thyroid gland is malfunctioning (hypothyroidism). A goiter can also occur in a gland that is producing too much thyroid hormone (hyperthyroidism) or even the correct amount of hormone (euthyroidism). Goiter indicates there is a condition present which is causing the thyroid to grow abnormally.

Q55: Define fluoride deficiency and toxicity?

Answer

Fluoride Toxicity

Fluoride toxicity or fluoride poisoning is a condition in which more fluoride is taken than the amount required for normal growth, development, and metabolism. Fluoride toxicity is characterized by a variety of signs and symptoms. Poisoning most commonly occurs following ingestion of a conspicuous amount of fluoride-containing products. Symptoms or signs usually occur within minutes of exposure. Fluoride is formed in

many common household products toothpaste, insecticides etc.,

Fluoride deficiency

Fluoride deficiency result when the amount of its up taken is less than required. Fluoride deficiency results in Brittle bones or demineralization of bones. Fluoride deficiency can lead to a higher likelihood of developing bone fractures and possibly even steoporesis.

We can combine these two half equations to give the overall reaction as follows:



Q56. What happen when HI is added to conc. H_2SO_4 ?

Answer

Iodide ions are stronger reducing agents they are oxidized to I_2 to I_2 by the conc. H_2SO_4 . ($2\text{I}^{-1} \longrightarrow \text{I}_2 + 2\text{e}^{-}$).

The reduction of the sulphuric acid is more complicated than before. The iodide ions are powerful enough reducing agents to reduce it.


This reduction takes place in following steps:

- First to sulphur dioxide (oxidation state of S = +4).
- Then to sulphur itself (oxidation state of S = 0).
- Then formation of H_2S (oxidation state of S = -2).

The most important of this mixture of reduction products is probably the H_2S . The half-reaction for its formation is:



On combing these text two half equations




Quick Quiz

Q57. HI is stronger acid then HF why?

Answer

Due to very high electronegativity of F (EN = 4). HF is highly polar molecule. Due to this reason strong hydrogen bonds are formed ammg H-F molecules.




On the other hand $\text{I}^{\ominus}$ is a large sized atom which can exist more freely. So HI releases more hydrogen ions. That is why HI is stronger acid then HF.

Q58. Although hydrogen bonding in HF is stranger than in H_2O but H_2O has much higher boiling point then that of H_2O , Why?

Answer

HF molecule may form one hydrogen bond whereas one H_2O molecule may form two H-bonds.



That is why boiling point of H_2O (100°C) is higher than that of HF (98°C).

Q59. The acidic character of hydrides of group VII-A elements increases on descending the group, Why?

Answer

Atomic sizes of halogens increase from F to I in group VII-A as a result stability of halogen ions increases. That's why the ability of halo acid to release H^+ ion increases, from HF to HI.

Q60. Illustrate the oxidizing properties of halogens by giving examples of two typical reactions.

Answer

A substance which has high electronegativity and ability to accept one or more electrons is said to show oxidizing property. Also the electron affinity values of halogens are high. But the electron affinity decreases from F to I. So the oxidizing power of halogens also decreases from F to I.

Upper halogen in group VII-A may displace lower halogen that is F_2 may displace lower halogen that is Cl^- ions then Cl_2 may displace Br^- ions from their solution.



Q61. Give reason

i) F is better oxidizing agent than Cl

ii) Electronegativity of halogens decreases in the order $\text{F} > \text{Cl} > \text{Br} > \text{I}$

Answer (i)

F has the highest electronegativity ($\text{EN} = 4$) whereas Cl has electronegativity ($\text{EN} = 3.2$). Therefore F may attract the electrons with more force and may oxidize any substance more than that of Cl.

Answer (ii)

Atomic sizes of halogens increase from F to I due to increase in number of shells from F to I. As a result pull of nucleus to attract the shared pair of electrons decreases from F to I.

F	>	Cl	>	Br	>	I
F	>	Cl	>	Br	>	I
E.N = Value = 4.0		3.2		3.0		2.5

Q62. Give different steps to obtain iron from its ores.

Answer


Iron has the following ores:

Magnetite	Fe_3O_4
Haematite	Fe_2O_3
Limonite	$\text{Fe}_2\text{O}_3 \cdot 3\text{H}_2\text{O}$

Iron is mostly obtained from magnetite and hematite.

Steps to extract metal from their ores:

- i) **Removal of gangue or matrix:**
In this step useless material from ores is removed manually or mechanically.
- ii) **Crushing**
In this step large pieces are broken down into small pieces with the crushers.
- iii) **Grinding**
Teema Ball mill is oftenly used to grind the ores without contamination.
- iv) **Concentration (Magnetic concentration of Iron)**
Different methods are used to concentrate the ore before sending to furnace. In case of iron magnetic separation method is used.
In this method powdered iron ore is passed over a magnetic pulley using rubber conveyer belts.



VI) Refining of metals (Open hearth Process)

In this process metal is heated in a fire bricks lining furnace. Hot air is through the molten metal which oxidized and remove the impurities on the top of molten metal or skimmed off.

Q63. What is bleaching powder? Give its common applications.

Answer

Bleaching powder is a mixture of calcium hypo chloride and calcium chloride $\text{Ca}(\text{OCl})_2$ CaCl_2 , with some slaked lime ($\text{Ca}(\text{OH})_2$).

Application of bleaching powder


- i) **Removal of stains**
Bleaching powder is used to remove stain from clothe and different materials to eliminate mildew and other stubborn stains.
- ii) To Disinfect and sterilize the baby toys and other home applications.

- iii) Removal of weeds, moss and algae from gardens and walk ways.
- iv) Bleaching powder is used to keep freshness of cut flowers by adding it to the solutions in which flowers are placed as it kills bacteria.
- v) Disinfection of drinking water and swimming pool water.
- vi) Bleaching of sugar in sugar industry.

Q64. Give different uses of halogens.


Answer

- i) Manufacturing Teflon: (Polytetra fluoroethylene) i.e.



Teflon is used to store HF and other acids which corrode metals.

Flourine is used in extraction of uranium and their isotope. Separation UF_6 is manufactured to separate ^{238}U and ^{235}U .



Q65. Write notes an toxicity of fluorides and deficiency of fluorides.

Answer

Flouride Toxicity

Flouride toxicity or fluoride poisoning is a condition is which more fluoride is takes than the amount required for normal growth, development an metabolism. Flouride toxicity is characterized by a variety of signs and symptoms. Poisoning most commonly occurs following ingestion of conspicuous amount of flouride entraining products symptoms or set usually occurs within mixtures of exposure. Flouride is found in many common household products e.g. toothpaste, dietary supplementary insecticides rodenticide etc. fluoride toxicity results.

- 1) Arthrit
- 2) Stiff painful joints with or wistful swelling.
- 3) Asthma, especially after showering.
- 4) Painful bony lumps where tendons and ligaments attach to bones.

Q65. What problems are caused in human body by deficiency of Iodine?

Answer

As human body cannot synthesize iodine so it is an essential element. The deficiency of iodine in human body leads to a disease goiter.

Goiter

The term goiter refers to abnormal enlargement of thyroid glands due to deficiency of iodine in diet. It results in swelling in neck. The presence of goiter does not means that the thyroid gland is malfunctioning (hypothyroidism): A goiter can also occur in a gland that is producing too much thyroid ramones (hyperthyroidism).

I-A	II-A	III-A	IV-A
+1	+2	+3, 5	2+,4+,6

Trends within a Group


The oxidation state of group I-A and II-A elements is fixed. But in P-block when we move from top to bottom the atomic size increases as a result the ability of atoms to loose electrons also increases.

For example in group IV-A


C and Si show +4 oxidatin state. In case of Ge, Sn and Tb only two nP electrons from the ns^2p^2 configuration are lost. We get the elements in +2 oxidatin sates remain inert and hence are not lost in the formation of M^{2+} cations. This pair of ns^2 electrons is called inert pair of electrons. Since the group the stability of +2 oxidation state also increases from Ge^{2+} to Pb^{2+} i.e $Ge^{2+} < Sn^{2+} < Pb^{2+}$.

When all the four ns^2p^2 electrons are lost we get the element in +4 oxidation state, i.e. M^{4+} cations are formed. On descending the group stability of +4 oxidation state decreases i.e the stability of M^{4+} cations decreases from Ge^{4+} to Pb^{4+} i.e. $Ge^{4+} > Sn^{4+} > Pb^{4+}$.

Compounds of Ge^{2+} are less stable than those of Ge^{4+} and hence the compounds of Ge^{2+} are readily charged oxidised into those of Ge^{4+} act as strong reducing agents while those of Ge^{4+} act as oxidising agents.



On similar grounds it can be shown that the compounds of Sn^{2+} are less stable than those of Sn^{4+} into those of Sn^{4+} . In other words, compounds of Sn^{2+} act as strong reducing agents while those of Sn^{4+} act as oxidising agents.



When we compare the stability of the compounds of Pb^{2+} and Pb^{4+} ions, we find that Pb^{2+} than those of Pb^{4+} ($PbCl_4$) and hence the compounds of Pb^{4+} are readily charged (reduced) into those of Pb^{2+} . In other words compounds of Pb^{4+} act as strong oxidising agents while those of Pb^{2+} act as reducing agents.

